

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)

2. REPORT TYPE
Technical Papers

3. DATES COVERED (From - To)

4. TITLE AND SUBTITLE

5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S)

5d. PROJECT NUMBER

2303

5e. TASK NUMBER

M208

5f. WORK UNIT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

8. PERFORMING ORGANIZATION
REPORT

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

10. SPONSOR/MONITOR'S
ACRONYM(S)

11. SPONSOR/MONITOR'S
NUMBER(S)

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

1121 013

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:

a. REPORT

Unclassified

b. ABSTRACT

Unclassified

c. THIS PAGE

Unclassified

17. LIMITATION
OF ABSTRACT

A

18. NUMBER
OF PAGES

19a. NAME OF RESPONSIBLE
PERSON

Leilani Richardson

19b. TELEPHONE NUMBER

(include area code)

(661) 275-5015

Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. Z39.18

62

separate items are enclosed

MEMORANDUM FOR PR (In-House Publication)

FROM: PROI (TI) (STINFO)

30 Jun 2000

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2000-144**
M. Fajardo, S. Tam, "High Resolution Infrared Absorption Spectroscopy in Doped Parahydrogen
Solids: CO/pH₂ – a Molecular Thermometer"

3rd International Conference on Cryocrystals and Quantum Crystals (Statement A)
(Szkarska Poreba, Poland, 28 Jul – 04 Aug 00) (Submission Deadline: 28 Jul 00)

1. This request has been reviewed by the Foreign Disclosure Office for: a.) appropriateness of distribution statement, b.) military/national critical technology, c.) export controls or distribution restrictions, d.) appropriateness for release to a foreign nation, and e.) technical sensitivity and/or economic sensitivity.

Comments: _____

Signature _____ Date _____

2. This request has been reviewed by the Public Affairs Office for: a.) appropriateness for public release and/or b) possible higher headquarters review.

Comments: _____

Signature _____ Date _____

3. This request has been reviewed by the STINFO for: a.) changes if approved as amended, b.) appropriateness of distribution statement, c.) military/national critical technology, d.) economic sensitivity, e.) parallel review completed if required, and f.) format and completion of meeting clearance form if required

Comments: _____

Signature _____ Date _____

4. This request has been reviewed by PR for: a.) technical accuracy, b.) appropriateness for audience, c.) appropriateness of distribution statement, d.) technical sensitivity and economic sensitivity, e.) military/national critical technology, and f.) data rights and patentability

Comments: _____

APPROVED/APPROVED AS AMENDED/DISAPPROVED

LESLIE S. PERKINS, Ph.D (Date)
Staff Scientist
Propulsion Directorate

High Resolution Infrared Absorption Spectroscopy in Doped Parahydrogen Solids: CO/pH₂ -- a Molecular Thermometer

Mario E. Fajardo, and Simon Tam

USAF Research Laboratory, AFRL/PRSP, Bldg. 8451, Edwards AFB, CA 93524-7680
mario_fajardo@ple.af.mil

20021121 013

- * HEDM Cryosolid Propellants
- * Trapping of Metal Atoms in Cryogenic Solid Hydrogen
- * Rapid Vapor Deposition of Transparent Parahydrogen (pH₂) Solids
- * High Resolution IR Absorption Spectroscopy in Doped pH₂ Solids
- * CO/pH₂ "Thermometer" Depositions
- * Summary

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

HEDM Cryosolid Propellants Payoffs

Increased Specific Impulse

$$I_{sp} \propto \sqrt{\Delta H_{sp}}$$

$$\text{LOX/LH}_2 : I_{sp} = 390 \text{ s}$$

$$5\% \text{ B/sH}_2 + \text{LOX} : I_{sp} = 500 \text{ s (+30\%)*}$$

* calculated for $P_{\text{chamber}} = 1000 \text{ PSIA}$, $P_{\text{exhaust}} = 14.7 \text{ PSIA}$

Greater Propellant Density

liquid H_2 @ 20 K : $\rho = 0.070 \text{ g/cm}^3$

solid H_2 @ 2 K : $\rho = 0.087 \text{ g/cm}^3$ (+25%)

50/50 liquid He/solid H_2 : $\rho = 0.105 \text{ g/cm}^3$ (+50%)


Dopant recombination/reaction in solid pH_2

* ideally:




isolated atoms


* in practice:



recombination




reaction




both

Experimental Diagram (c1993)



Transmission Spectrum of Li/nH_2 , $d \approx 10 \mu$



M.E. Fajardo, J. Chem. Phys. **98**, 110 (1993).

Optical Scattering in Solid Hydrogen

Crystal Growing and Quality (p. 81)

“There is a considerable art to growing hydrogen crystals of high quality. Good crystals are always grown slowly from the melt; a rapid freeze from the gas produces snow.”


Crystallite Light Scattering (p. 83)

“The reason that a good hydrogen crystal is so hard to see is its low refractive index...an estimated 1.16!


Yet a 1 mm-thick layer of hydrogen crystallites can be a completely opaque brown-black.”

[P.C. Souers, Hydrogen Properties for Fusion Energy (UC Press, Berkeley, 1986)]

ortho- and para-hydrogen




Experimental Diagram (c1997)



M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).
S. Tam and M.E. Fajardo, Rev. Sci. Instrum. **70**, 1926 (1999).


Transmission Spectrum of B/pH₂ d_≈1mm



M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).

S. Tam, M. MacIer, M.E. DeRose, and M.E. Fajardo, J. Chem. Phys., submitted (2000).

IR Absorption of 6 mm Thick pH_2 Solid




Non-observation of the $Q_1(0)$ transition demonstrates the absence of oH_2 impurities, and that the microscopic structure is not amorphous or porous.

Observation of $S_1(0)$ transition demonstrates the absence of inversion symmetry for some H_2 molecular environments.

[van Kranendonk and Gush, Phys. Lett. **1**, 22 (1962)]

M.E. Fajardo and S. Tam, J. Chem. Phys. **108**, 4237 (1998).

Raman Spectra of pH_2 Solids



Mixed hcp/fcc as-deposited structure, anneals to hcp.

[G.W. Collins, et al., Phys. Rev. B **53**, 102 (1996)]

(d) sample in (c) warmed to 4.5 K.
(c) 4.5 mm sample as deposited at 3.3 K ($\Phi = 290$ mmol/hr).

(b) sample in (a) warmed to 4.5 K.
(a) 6 mm sample as deposited at 3.1 K ($\Phi = 200$ mmol/hr).

High Res. IR Spectroscopy in Solid pH_2






FIG. 1. Apparatus for the simultaneous spectroscopy of the infrared and Raman transitions. The nonlinearity of LiNbO_3 is used for the former and that of solid H_2 is used for the latter. D.M., dichroic mirror; S. A., spectrum analyzer; P. B., polarizer beamsplitter.

ν_4 CH₄/pH₂ IR Absorptions (res = 0.01 cm⁻¹)




S. Tam, M.E. Fajardo, H. Katsuki, H. Hoshina, T. Wakabayashi, and T. Momose, J. Chem. Phys. **111**, 4191 (1999).

ν_4 CH₄/pH₂ IR Absorptions




- (a) Rapid Vapor Deposited sample: as-deposited at 2.4 K
- (b) Rapid Vapor Deposited sample: annealed to 4.8 K
- (c) Enclosed Cell Condensed sample: cooled to 4.8 K

V_4 CH₄/pH₂ Energy Levels




S. Tam, M.E. Fajardo, H. Katsuki, H. Hoshina, T. Wakabayashi, and T. Momose, J. Chem. Phys. **111**, 4191 (1999).

CO_2/pH_2 Trapping Sites




S. Tam and M.E. Fajardo, Fiz. Nizk. Temp. [Low Temp. Phys.] accepted (2000).

CO₂/pH₂ IR Absorptions (res = 0.008 cm⁻¹)

as-deposited at $T = 2.4 \text{ K}$

(c) 1.2 ppm CO₂/pH₂
(b) 0.04 ppm CO₂/pH₂
(a) 0.01 ppm CO₂/pH₂

$^{13}\text{C}^{16}\text{O}_2/\text{pH}_2$ vs. $^{12}\text{C}^{16}\text{O}_2/\text{pH}_2$




annealed, $T = 2.4 \text{ K}$

$d = 1.0 \text{ mm}$

2 ppm $^{13}\text{C}^{16}\text{O}_2$ and
2 ppm $^{12}\text{C}^{16}\text{O}_2$

88 PPM HCl/pH₂



Gas Phase $(\text{HCl})_2$

High resolution, jet-cooled infrared spectroscopy of $(\text{HCl})_2$: Analysis of ν_1 and ν_2 HCl stretching fundamentals, interconversion tunneling, and mode-specific predissociation lifetimes

Michael D. Schuder,^{a)} Christopher M. Lovejoy,^{b)} Robert Lascola,^{c)} and David J. Nesbitt^{d)}
*Joint Institute for Laboratory Astrophysics, National Institute of Standards and Technology and
 University of Colorado, and the Department of Chemistry and Biochemistry, University of Colorado,
 Boulder, Colorado 80309*

(Received 5 April 1993; accepted 7 June 1993)






FIG. 1. Vibrationally averaged structure and internal coordinates for HCl dimer. The intermolecular axis R connects the HCl centers of mass. The internal angles, θ_1 and θ_2 , are measured from the intermolecular axis to the HCl bonds r . The torsion angle, $\phi = \phi_1 - \phi_2$, is shown at 180° (planar). The minimum energy configuration shown is for $\theta_1 = 16^\circ$, $\theta_2 = 87^\circ$ with $\phi_1 - \phi_2 = 180^\circ$. The HCl subunit on the left is referred to as the bonded HCl with an associated vibration labeled ν_2 . The proton on the other HCl is not involved with the hydrogen bond, and this subunit is referred to as the free HCl , with a vibration labeled ν_1 .




$(\text{HCl})_2/\text{pH}_2$ isotopomers




analysis in collaboration with D.T. Anderson, U. Wyoming.

Co-operative IR absorptions




analysis in collaboration with R.J. Hinde, U. Tennessee, Knoxville.

80 PPM CO/pH₂ (res = 0.1 cm⁻¹)



analysis in collaboration with T. Momose, Kyoto U.

$(\text{CO})_n/\text{pH}_2$ IR Absorptions




$\approx 6 \text{ }\mu\text{mol CO}$ in
each sample

$d = 1.7 \text{ mm}$





$d = 20 \text{ }\mu\text{m}$

$d = 4.7 \text{ }\mu\text{m}$

Experimental Diagram



Experimental Protocol



			
pure PbS	CO_2/PbS	CO_2/PbS	pure PbS

thermometer
experiments:


control
experiment

100
200
300
400
500
600
700
800
900
1000

CO₂/pH₂ Thermometer Peak



Calibration of CO/pH₂ Thermometer



simplified Boltzmann:

$$A_{2135} = N \exp(-E/T_{CO})$$


$$N = 0.0860$$

$$E = 7.896 \text{ K}$$


include $Q_{rot} \Rightarrow E \approx 11 \text{ K}$

$$6 B_{CO/pH_2} \approx 12 \text{ K}$$


Deposition Temperatures



Cryostat Thermal Response



Deposition Heat Loads



heat fluxes calculated from
 $\Delta T = T_B - T_B(\text{prior dep.})$

slope of fit line \Rightarrow
 $E_{\text{dep}} = 3.3 \text{ kJ/mol}$

compare with:
 $E_{\text{vap}} = 1.1 \text{ kJ/mol}$

\leftarrow note post-deposition
radiative(?) heat loads

pH₂ Thermal Conductivity Calculations

expt.#	$\Delta x(\text{cm})$	$\Delta T_{\text{max}}(\text{K})$	$\Delta T_{\text{min}}(\text{K})$	$(\dot{Q}/A)_{\text{max}}$	$(\dot{Q}/A)_{\text{min}}$	K_{min}	K_{max}
1	0.108	0.61	0.21	9.2	6.9	1.2	4.7
2	0.203	1.19	0.79	13.8	10.7	1.8	3.5
3	0.232	1.74	1.34	16.6	12.5	1.7	2.9

Units: \dot{Q}/A (mW/cm²), κ (mW/cm-K).

Summary: $\kappa = 3(\pm 2)$ mW/cm-K, averaged over $2 < T < 5$ K range.

Comparison with Literature

Previous studies on pH_2 solids grown from the gas phase in an enclosed cell near 10 K

$$\Rightarrow \bar{\kappa} \approx 4000 \text{ mW/cm-K}$$

$$\& L_{ph} \sim 0.1 \text{ cm.}$$

$$\kappa = C v L_{ph} / 3$$

[V.G. Manzhelii, B.Ya. Gorodilov, and A.I. Krivchikov, Low Temp. Phys. 22, 131 (1996)]




FIG. 1. Thermal conductivity of crystals of pure pH_2 and pH_2 with Ne impurity (the concentration in ppm are indicated); the solid lines are calculated results.

Suggests $L_{ph} \sim 1 \mu m$ in our rapid vapor deposited solids.

Summary

- * Demonstrated production of millimeters-thick transparent pH_2 solids by rapid vapor deposition.
- * Demonstrated that vapor deposited pH_2 solids are densest close-packed solids, NOT amorphous.
- * Demonstrated suitability of vapor deposited pH_2 solids as hosts for high resolution IR absorption spectroscopy of chemically interesting dopants.
- * Generalized phenomenon of dopant induced IR activity.
- * Exploited CO/pH_2 spectroscopy to probe sample temperature during deposition, and to estimate thermal conductivity.

Collaborators

- * Mr. Simon Tam and Ms. Michelle E. DeRose, AFRL/PRSP
responsible for our experimental data.
- * Prof. Takamasa Momose, Kyoto U.
spectroscopy of CH_4 , C_{60} , and CO doped pH_2 solids.
- * Prof. Robert J. Hinde, U. Tennessee at Knoxville
dopant-induced IR absorptions.
- * Prof. David T. Anderson, U. Wyoming
spectroscopy of $(\text{HCl})_2$ in solid pH_2 .